arXiv:math/0608787v3 [math.CA] 12 Jul 2007

AN APPLICATION OF λ-METHOD ON INEQUALITIES OF SHAFER-FINK'S TYPE

Branko J. Malešević

Abstract. In this article λ-method of Mitrinović-Vasić [1] is applied to improve the upper bound for the arc sin function of L. Zhu [4].

1. Inequalities of Shafer-Fink's type

D. S. Mitrinović in [1] considered the lower bound of the arc sin function, which belongs to R. E. Shafer. Namely, the following statement is true.

Theorem 1.1 *For* $0 \leq x \leq 1$ *the following inequalities are true*:

$$\frac{3x}{2+\sqrt{1-x^2}} \leq \frac{6(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}} \leq \arcsin x\,. \tag{1}$$

A. M. Fink proved the following statement in [2] .

Theorem 1.2 *For* $0 \leq x \leq 1$ *the following inequalities are true*:

$$\frac{3x}{2+\sqrt{1-x^2}} \leq \arcsin x \leq \frac{\pi x}{2+\sqrt{1-x^2}}. \tag{2}$$

B. J. Malešević proved the following statement in [3].

Theorem 1.3 *For* $0 \leq x \leq 1$ *the following inequalities are true*:

$$\frac{3x}{2+\sqrt{1-x^2}} \leq \arcsin x \leq \frac{\dfrac{\pi}{\pi-2}x}{\dfrac{2}{\pi-2}+\sqrt{1-x^2}} \leq \frac{\pi x}{2+\sqrt{1-x^2}}. \tag{3}$$

The main result of the article [3] can be formulated with the next statement.

Proposition 1.4 *In the family of the functions*:

$$f_b(x) = \frac{(b+1)x}{b+\sqrt{1-x^2}} \quad (0 \leq x \leq 1), \tag{4}$$

according to the parameter $b > 0$, *the function* $f_2(x)$ *is the greatest lower bound of the* $\arcsin x$ *function and the function* $f_{2/(\pi-2)}(x)$ *is the least upper bound of the* $\arcsin x$ *function.*

Research partially supported by the MNTRS, Serbia, Grant No. 144020.
Keywords: Inverse sine; Upper and lower bounds; Shafer-Fink type inequality.

L. Zhu proved the following statement in [4].

Theorem 1.5 *For $x \in [0,1]$ the following inequalities are true*:

$$\begin{aligned}\frac{3x}{2+\sqrt{1-x^2}} &\leq \frac{6(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}} \leq \arcsin x \\ &\leq \frac{\pi(\sqrt{2}+\frac{1}{2})(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}} \leq \frac{\pi x}{2+\sqrt{1-x^2}}.\end{aligned} \tag{5}$$

In this article we further improve the upper bound of the arc sin function. Namely, in the next section we will give proof of the following theorem:

Theorem 1.6 *For $x \in [0,1]$ the following inequalities are true*:

$$\begin{aligned}\frac{3x}{2+\sqrt{1-x^2}} &\leq \frac{6(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}} \leq \arcsin x \\ &\leq \frac{\frac{\pi(2-\sqrt{2})}{\pi-2\sqrt{2}}(\sqrt{1+x}-\sqrt{1-x})}{\frac{\sqrt{2}(4-\pi)}{\pi-2\sqrt{2}}+\sqrt{1+x}+\sqrt{1-x}} \\ &\leq \frac{\pi(\sqrt{2}+\frac{1}{2})(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}} \leq \frac{\pi x}{2+\sqrt{1-x^2}}.\end{aligned} \tag{6}$$

Remark 1.7 *Using numerical method from* [5] *we have the following conclusions*:

$\mathbf{1^0}$. *For values $x \in (0, 0.387\,266\,274\ldots)$ the following inequality is true*:

$$\arcsin x < \frac{\frac{\pi}{\pi-2}x}{\frac{2}{\pi-2}+\sqrt{1-x^2}} < \frac{\pi(\sqrt{2}+\frac{1}{2})(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}}, \tag{7}$$

and for values $x \in (0.387\,266\,274\ldots, 1)$ the following inequality is true:

$$\arcsin x < \frac{\pi(\sqrt{2}+\frac{1}{2})(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}} < \frac{\frac{\pi}{\pi-2}x}{\frac{2}{\pi-2}+\sqrt{1-x^2}}. \tag{8}$$

Numerically determined constant $c = 0.387\,266\,274\ldots$ is the unique number where the previous bounds have the same values over $(0,1)$.

$\mathbf{2^0}$. *For values $x \in (0,1)$ the following inequality is true*:

$$\arcsin x < \frac{\frac{\pi(2-\sqrt{2})}{\pi-2\sqrt{2}}(\sqrt{1+x}-\sqrt{1-x})}{\frac{\sqrt{2}(4-\pi)}{\pi-2\sqrt{2}}+\sqrt{1+x}+\sqrt{1-x}} < \frac{\frac{\pi}{\pi-2}x}{\frac{2}{\pi-2}+\sqrt{1-x^2}}. \tag{9}$$

2. The main results

In this article, using λ-method of Mitrinović-Vasić we give an analogous statement to Proposition 1.4. Let us notice that from inequality given by L. Zhu [4]:

$$\frac{6(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}} \leq \arcsin x \leq \frac{\pi(\sqrt{2}+\frac{1}{2})(\sqrt{1+x}-\sqrt{1-x})}{4+\sqrt{1+x}+\sqrt{1-x}}, \tag{10}$$

for $x \in [0,1]$, we can conclude that the function $\varphi(x) = \arcsin x$ has a lower bound and upper bound in the family of the functions:

$$\Phi_{\alpha,\beta}(x) = \frac{\alpha(\sqrt{1+x}-\sqrt{1-x})}{\beta+\sqrt{1+x}+\sqrt{1-x}} \quad (0 \leq x \leq 1), \tag{11}$$

for some values of parameters $\alpha, \beta > 0$. Next for $x = 0$ it is true that $\Phi_{\alpha,\beta}(0) = 0$, for $\alpha, \beta > 0$. On the other hand, for values $x \in (0,1]$ it is true:

$$\Phi_{\alpha_1,\beta_1}(x) > \Phi_{\alpha_2,\beta_2}(x) \Longleftrightarrow \alpha_1\beta_2 - \alpha_2\beta_1 > (\alpha_2-\alpha_1)(\sqrt{1+x}+\sqrt{1-x}), \tag{12}$$

for $\alpha_{1,2}, \beta_{1,2} > 0$. Let us apply λ-method of Mitrinović-Vasić on the considered two-parameters family $\Phi_{\alpha,\beta}(x)$ in order to determine the bounds of the function $\varphi(x)$ under the following conditions:

$$\Phi_{\alpha,\beta}(0) = \varphi(0) \quad \text{and} \quad \frac{d}{dx}\Phi_{\alpha,\beta}(0) = \frac{d}{dx}\varphi(0). \tag{13}$$

It follows that $\alpha = \beta + 2$. In that way we get one-parameter subfamily:

$$f_\beta(x) = \Phi_{\beta+2,\beta}(x) = \frac{(\beta+2)(\sqrt{1+x}-\sqrt{1-x})}{\beta+\sqrt{1+x}+\sqrt{1-x}} \qquad (0 \leq x \leq 1), \tag{14}$$

according to the parameter $\beta > 0$. For that family the condition (13) is true:

$$f_\beta(0) = \varphi(0) \;\text{ and }\; \frac{d}{dx}f_\beta(0) = \frac{d}{dx}\varphi(0). \tag{15}$$

Additionally, we have:

$$\frac{d^2}{dx^2}f_\beta(0) = \frac{d^2}{dx^2}\varphi(0) \;\text{ and }\; \frac{d^3}{dx^3}f_\beta(0) = \frac{d^3}{dx^3}\varphi(0) + \frac{4-\beta}{4(2+\beta)} \tag{16}$$

and

$$\frac{d^4}{dx^4}f_\beta(0) = \frac{d^4}{dx^4}\varphi(0) \;\text{ and }\; \frac{d^5}{dx^5}f_\beta(0) = \frac{d^5}{dx^5}\varphi(0) + \frac{3(128+18\beta-13\beta^2)}{16(2+\beta)^2}. \tag{17}$$

Let us notice that for the family of the functions $f_\beta(x)$, on the basis of (12), for values $x \in (0,1]$ the following equivalence is true:

$$f_{\beta_1}(x) > f_{\beta_2}(x) \Longleftrightarrow \beta_1 < \beta_2, \tag{18}$$

for $\beta_{1,2} > 0$. Let us emphasize that there is a better upper bound $f_{b_1}(x)$ than upper bound $\Phi_{\pi(\sqrt{2}+1/2),4}(x)$ of the function $\varphi(x)$ over $(0,1]$. It is true that the parameter $\beta = b_1$ fulfils:

$$f_{b_1}(1) = \varphi(1) = \frac{\pi}{2}, \tag{19}$$

hence:

$$b_1 = \frac{\sqrt{2}(4-\pi)}{\pi - 2\sqrt{2}} = 3.876\,452\,527\ldots < 4. \tag{20}$$

Let us prove that the function $f_{b_1}(x)$ is the upper bound of the function $\varphi(x)$ over $[0,1]$. Let us define the function:

$$h(x) = f_{b_1}(x) - \varphi(x) \tag{21}$$

for $0 \leq x \leq 1$. For the function $h(x)$ we introduce two substitutions $x = \cos t$ $\big(t \in [0, \frac{\pi}{2}]\big)$ and $t = 4 \operatorname{arc\,tg} u$ $\big(u \in [0, \operatorname{tg}\frac{\pi}{8}]\big)$ respectively, and we get a new function:

$$\mathrm{w}(u) = h\big(\cos(4 \operatorname{arc\,tg} u)\big) = \frac{\sqrt{2}(b_1+2)(u^2+2u-1)}{(\sqrt{2}-b_1)u^2 - 2\sqrt{2}u - b_1 - \sqrt{2}} - \frac{\pi}{2} + 4 \operatorname{arc\,tg} u \tag{22}$$

for $0 \leq u \leq \operatorname{tg}\frac{\pi}{8} = \sqrt{2}-1$. Then:

$$\begin{aligned} \frac{d}{du}\mathrm{w}(u) \;=\; & \Big(\big(4b_1^2+2\sqrt{2}b_1^2-8b_1-4\sqrt{2}b_1-8\big)u^4+\big(-4\sqrt{2}b_1^2+8\sqrt{2}b_1-32\big)u^3+ \\ & \big(8b_1^2-16b_1-16\big)u^2+\big(-4\sqrt{2}b_1^2+8\sqrt{2}b_1+32\big)u+ \\ & \big(4b_1^2-2\sqrt{2}b_1^2-8b_1+4\sqrt{2}b_1-8\big)\Big)\Big/ \\ & \Big(\big(u^2+1\big)\big(b_1u^2-\sqrt{2}u^2+2\sqrt{2}u+b_1+\sqrt{2}\big)^2\Big). \end{aligned} \tag{23}$$

All solutions of the equation $\frac{d}{du}\mathrm{w}(u) = 0$ are determined by terms:

$$\begin{aligned} u_{1,4} &= \frac{2\sqrt{2} \mp \sqrt{-b_1^4+4b_1^3+4b_1^2-16b_1}}{b_1^2-2b_1+2\sqrt{2}-4}\,, \\ u_{2,3} &= \sqrt{2}-1\,; \end{aligned} \tag{24}$$

or by numerical values: $u_1 = 0.0869\ldots$, $u_{2,3} = 0.4142\ldots$, $u_4 = 0.8400\ldots$. The function $\mathrm{w}(u)$ has local maximum at the point u_1 and $\mathrm{w}(0) = \mathrm{w}(\sqrt{2}-1) = 0$. Hence $\mathrm{w}(u) \geq 0$ for $u \in [0, \sqrt{2}-1]$. Therefore the function:

$$f_{b_1}(x) = \frac{\dfrac{\pi(2-\sqrt{2})}{\pi-2\sqrt{2}}\big(\sqrt{1+x}-\sqrt{1-x}\big)}{\dfrac{\sqrt{2}(4-\pi)}{\pi-2\sqrt{2}} + \sqrt{1+x} + \sqrt{1-x}} \tag{25}$$

is the upper bound of $\varphi(x)$ over $[0,1]$. Let us notice that, for values $x \in (0,1]$, on the basis (12), the following inequalities are true:

$$\varphi(x) < f_{b_1}(x) = \Phi_{b_1+2,b_1}(x) < \Phi_{\pi(\sqrt{2}+1/2),4}(x). \tag{26}$$

Let us prove that the function $f_{b_1}(x)$ is the least upper bound of the function $\varphi(x)$ from the family (14). The following implication is true:

$$b_1 < b \implies f_b(1) < f_{b_1}(1) = \varphi(1) = \frac{\pi}{2}. \tag{27}$$

Hence for $b > b_1$ the function $f_b(x)$ is not the upper bound for the function $\varphi(x)$ over $[0,1]$. According to the previous consideration we can conclude that the function $f_{b_1}(x)$ is the least upper bound of the function $\varphi(x)$ over $[0,1]$.

The lower bound of the function $f_4(x)$ of the function $\varphi(x)$ over $[0,1]$, which belongs to R. E. Shafer, according to formulas (15) - (17), has at $x = 0$ the root of the fifth order. Let us prove that the function $f_4(x)$ is the greatest lower bound of the function $\varphi(x)$ from the family (14). For fixed $b \in (b_1, 4)$ let us define the function:

$$g(x) = \begin{cases} \alpha & : \quad x = 0, \\ \dfrac{f_b(x) - \varphi(x)}{x^3} & : \quad x \in (0,1]; \end{cases} \tag{28}$$

with the constant:

$$\alpha = \frac{\dfrac{d^3}{dx^3} f_b(0) - \dfrac{d^3}{dx^3}\varphi(0)}{6} = \frac{4-b}{24\big(2+b\big)} > 0. \tag{29}$$

The function $g(x)$ is continuous over $[0,1]$ and the following is true:

$$g(0) > 0 \quad \text{and} \quad g(1) < 0. \tag{30}$$

Therefore we can conclude that there is $c_b \in (0,1)$ such that $g(c_b) = 0$. Let us notice that $g(0) > 0$ and $g(c_b) = 0$. Then, there is some point $\xi_b \in (0, c_b)$ such that $g(\xi_b) > 0 \ \big(g \in \mathrm{C}[0, c_b]\big)$. This is sufficient for conclusion that, for each $b \in (b_1, 4)$, the function $f_b(x)$ is not the lower bound of the function $\varphi(x)$ over $[0,1]$. According to the previous consideration we can conclude that the function $f_4(x)$ is the greatest lower bound of the function $\varphi(x)$ over $[0,1]$.

On the basis of the previous consideration the following statement is true.

Proposition 2.1 *In the family of the functions*:

$$f_b(x) = \Phi_{b+2,b}(x) = \frac{(b+2)(\sqrt{1+x} - \sqrt{1-x})}{b + \sqrt{1+x} + \sqrt{1-x}} \qquad (0 \le x \le 1), \tag{31}$$

according to the parameter $b > 0$, the function $f_4(x)$ is the greatest lower bound of the arc sin x function and the function $f_{\sqrt{2}(4-\pi)/(\pi-2\sqrt{2})}(x)$ is the least upper bound of the arc sin x function.

Remark 2.2 *Let us emphasize that Theorem* 1.6 *has been recently considered in* [6] *and* [7]*. In the article* [7] *a simple proof of Theorem* 1.6 *based on "L'Hospital rule for monotonicity" is obtained.*

REFERENCES

[1] D. S. Mitrinović, P. M. Vasić: *Analytic inequalities*. Springer–Verlag 1970.

[2] A. M. Fink: *Two inequalities*, Univ. Beograd. Publ. Elektrotehn. Fak., Ser. Mat. **6** (1995), 48–49. (`http://pefmath.etf.bg.ac.yu/`)

[3] B. J. Malešević: *An application of λ-method on Shafer-Fink's inequality*, Univ. Beograd. Publ. Elektrotehn. Fak., Ser. Mat. 8 (1997), 90–92. (`http://pefmath.etf.bg.ac.yu/`)

[4] L. Zhu: *On Shafer-Fink inequalities*, Mathematical Inequalities & Applications, Vol. 8, Number 4, (2005), 571–574. (`http://www.mia-journal.com/`)

[5] B. J. Malešević: *One method for prooving inequalities by computer*, Journal of Inequalities and Applications, Vol. 2007, Article ID 78691, 8 pages, 2007. doi:10.1155/2007/78691. (`http://www.hindawi.com/journals/jia/`)

[6] L. Zhu: *A solution of a problem of Oppeheim*, Mathematical Inequalities & Applications, Vol. 10, Number 1, (2007), 57-61. (`http://www.mia-journal.com/`)

[7] L. Zhu: *On Shafer-Fink Type Inequality*, Accepted in Journal of Inequalities and Applications, Vol. 2007, ID 67430.
(`http://www.hindawi.com/journals/jia/`)

University of Belgrade,
Faculty of Electrical Engineering,
P.O.Box 35-54, 11120 Belgrade, Serbia
malesevic@etf.bg.ac.yu, **malesh@eunet.yu**

(Received 09/30/2006)
(Revised 05/08/2007)